


Instructions for use

EURneffy® 2 mg nasal spray, solution in single-dose container


adrenaline

Before you need to use it, fully familiarise yourself with EURneffy, including when and how it should be used.

EURneffy 2 mg nasal spray





Follow these instructions only when ready to use.

A  **Remove EURneffy 2 mg nasal spray from the packaging.**
Pull open the packaging to remove the EURneffy 2 mg nasal spray.

B  **Hold the nasal spray as shown.**
Hold the nasal spray with your thumb on the bottom of the plunger and one finger on either side of the nozzle.

- **Do not pull or push on the plunger.**
- **Do not test or pre-spray; each nasal spray has only one dose.**

C  **Insert the tip of nasal spray into a nostril until your fingers touch your nose.**
Keep the nozzle straight into the nose pointed toward your forehead. Do not angle the nasal spray to the inner or outer walls of the nose.


D  **Press the plunger up firmly until it snaps up and sprays into the nostril.**

  **Do not angle the nasal spray to the inner or outer walls of the nose.**

 **Seek emergency medical assistance**

Seek emergency medical assistance immediately (dial 999) as you might need further treatment for your anaphylaxis.

 **Monitor patient symptoms**

 If symptoms continue to worsen or reoccur after 5 minutes, or in case of any error in administration, use a new EURneffy nasal spray to give a second dose in the SAME nostril as the first dose and seek emergency medical assistance.

You should lie down with feet raised. If this makes you breathless, you can gently sit up. You should not change position suddenly and should lie down again as soon as possible. You should not stand up even if someone encourages you to. You should stay lying down even if you feel better. Pregnant women should lie on their left side. Unconscious patients should be placed on their side in the recovery position to prevent choking. If symptoms do not resolve, you should, if possible, remain with another person until medical assistance arrives.